

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

V. An Account of the Depth of Rain fallen from April 1, 1722, to April 1, 1723. Observed at Widdrington in Northumberland, and communicated to the Publisher by the Reverend Mr. Horsley.

Have kept an exact Account of what Rain has fallen the last Year in this Place, which I have here fent you. The weighing the Water and reducing it from Weight to Depth seem'd pretty troublesome, even when done in the easiest Method: To remedy this Inconvenience (besides a Funnel and proper Receptacle for the Rain) I use a cylindrical Measure and Gage. The Funnel is 30 Inches diameter, and the cylindrical Measure exactly 3; the Depth of the Measure is 10 Inches, and the Gage of the same Length, with each Inch divided into 10 equal Parts; or, instead of a Gage, the Inches and Divisions may be mark'd on the Side of the cylindrical Measure. The Apparatus is simple and plain, and it is easy to apprehend the Defign and Reason of the Contrivance; for the Diameter of the cylindrical Measure being just  $\frac{1}{10}$  of that of the Funnel, and the Measure exactly 10 Inches deep, 'tis plain that 10 Measures of Rain make an Inch in Depth; one Measure,  $\frac{1}{10}$ ; one Inch on the Gage,  $\frac{1}{100}$ ; and  $\frac{1}{10}$  of an Inch on the Gage,  $\frac{1}{1000}$ , &c. By this Means the Depth of any particular Quantity which falls, may be fet down with Ease and Exactness, and the

the whole at the End of each Month, or every Year,

may be fumm'd up without any Trouble.

By the following Account you'll see, that some of the Summer Months, particularly May and July, were very wet, and some of the Winter ones very dry; so that, one with another, this Year's Rain, as far as can well be conjectur'd, may be look'd upon as a Medium. And if so, it differs not above two or three Inches, from the mean Quantity of Rain which falls at Upminster, Paris, and Lisle, being less than at Lisle, and more than at the other two Places.

An Account of the Depth of Rain fallen from April 1, 1722, to April 1, 1723.

	Inch.
In April	1,015
In May	3,532
In June	2,570
In July	4,350
In August $    -$	2,132
In September	1,155
In October	,600
In November	2,205
In December	1,780
In Fanuary	1,225
In February	,485
In March	,195
In the whole Year	21,244